

FIGURE 1A




FIGURE 1B





FIGURE 1C






FIGURE 1D




FIGURE 1E




FIGURE 2A




FIGURE 2B




FIGURE 2C

4/31




FIGURE 2D




FIGURE 2E




FIGURE 3A




FIGURE 3B




FIGURE 3C






FIGURE 3D

7/31



8/31




FIGURE 4C




FIGURE 4D

9/31

FIGURE 5A





FIGURE 5B



10/31

FIGURE 6A






FIGURE 6B

FIGURE 6C

FIGURE 7A





FIGURE 7B



12/31

FIGURE 8




FIGURE 9A






FIGURE 9B





14/31

FIGURE 10



15/31



16/31

FIGURE 12




FIGURE 13




FIGURE 14A






FIGURE 14B



19/31

FIGURE 15

luciferase Activity in CPS/ μ g protein



20/31

FIGURE 16A




FIGURE 16B





FIGURE 17



22/31

FIGURE 18

Expression of SeAP 48h after cotransfection of NIH3T3s
using 6 nmol of RPR120535B/ μ g of DNA

In the presence of serum





FIGURE 19: Summarizing table of inhibition experiments with antisense transcripts of different lengths

Percentage inhibition of SeAP activity								
	size (pb)	antisense (pb)	Conserved sequence	Exp. 1	Exp. 2	Exp. 3	Exp. 4	Exp. 5
pSeAPAS	3447	1527						
pGJA1	2045	125	3' end of SeAP	70	85	94	90	92
pGJA2	1955	35	5' end of SeAP	24	52	63		
pGJA3	2123	203	5' end of SeAP	14	0	8		
pGJA9	2248	125 +203	2 ends	72	76	77	50	71


FIGURE 20

Evaluation of the control of the expression of SeAP with the combination of the SeAPantisense gene and the Tet Off system



25/31

FIGURE 21
Evaluation of the expression of SeAP 48h after transfection of NIH3T3
using 6 nmol RPR120535B / μ g DNA in the presence of serum



26/31

FIGURE 22





FIGURE 23 Regulation of the expression of SeAP 48h after transfection of NIH3T3s using 6 nmol IPR120535B/ μ g of DNA in the presence of serum



28/31

FIGURE 24 Decrease in background noise of the hPPARg2 system in C2C12 transfection, expression at day 5






FIGURE 25

NIH3T3 Transfection



30/31

FIGURE 26



ponasterone A

FIGURE 27

